

WISCONSIN FOREST TREE PLANTING HERBICIDES...2003

Table 3b. Timing of Herbicide Application for Commonly Planted Trees in Wisconsin. Wisconsin DNR, Forest Health Protection

Herbicide Trade Name	RED OAK	WHITE OAK	BLACK WALNUT	SUGAR MAPLE	SILVER MAPLE	WHITE ASH	GREEN ASH	HYBIRD POPLAR	RED PINE	WHITE PINE	JACK PINE	WHITE SPRUCE	NORWAY SPRUCE	WHITE CEDAR
Transline	Pre/Post		Pre/Post	Pre/Post		Pre/Post		Pre/Post	Pre/Post	Pre/Post		Pre/Post	Pre/Post	
Accord, Glypro, Glyphosate	Pre	Pre	Pre	Pre	Pre	Pre	Pre		Pre	Pre	Pre	Pre	Pre	Pre
Pronone 10G									Pre			Pre		
Velpar L									Pre			Pre		
Velpar DF									Pre			Pre		
Arsenal AC 1									Pre	Pre	Pre	Pre		
Linex 4L Lorox DF								Pre/Post						
Escort 4									Pre					
Pendulum	Pre/Post	Pre/Post	Pre/post	Pre/Post		Pre/Post			Pre/Post	Pre/Post		Pre/Post	Pre/Post	Pre/Post
Vantage	Pre/Post	Pre/Post	Pre/Post		Pre/Post	Pre/Post	Pre/Post	Pre/Post	Pre/Post	Pre/Post	Pre/Post	Pre/Post	Pre/Post	Pre/Post
Sim-Trol 90DF 2	Pre/Post		Pre/Post	Pre/Post					Pre/Post	Pre/Post		Pre/Post	Pre/Post	Pre/Post
Simazine 4L Simazine 90DF 2	Pre/Post			Pre/Post					Pre/Post	Pre/Post		Pre/Post	Pre/Post	Pre/Post
Oust XP 3	Pre/Post	Pre/Post				Pre	Pre/Post	Post		Post	Pre/Post	Post		

Note: "Pre" and "Post" refer to before and after tree planting.

FOOTNOTES:

1. Wait 3 to 6 months before planting. See supplemental labels.

2. The " Shelterbelts" section of the labels allows the use of these products in tree plantings for soil protection under a label interpretation. **SHAKE WELL** before pouring simazine 4L products from their containers and agitate simazine formulations in the spray tank to keep them well mixed.

3. Because occasional damage to trees has been observed at the higher application rates, the DNR recommendation for Oust applications is 0.5 - 1.0 oz. per acre on tree planting sites. The Oust label allows higher rates which vary depending on tree species. Tank mixes of Oust and simazine are being used successfully to control a wider range of weed species. Oust can be applied to the planting site in the fall before planting or in the

spring after planting; Oust should **not** be applied in the spring shortly before trees are planted.

4. Apply Escort the season before tree planting.

The information in these tables is dated. Label changes can make the information obsolete. Therefore the information in these tables should be used as a guide only and not considered recommendations.

To be safe always:

READ THE PACKAGE LABEL AND USE ACCORDINGLY.